

NSTEMI



GRACE _____	Troponin	Very high risk	High risk
	H0	O Hemodynamic instability O Cardiogenic shock O Recurrent/ongoing chest pain despite medical treatment O Life threatening arrhythmia O Mechanical complication O Acute HF related NSTEMI O Diffuse STD + STE aVR/V1	O Established Dx NSTEMI O Dynamic new ST-T change O Wellen, De Winter O Resuscitated cardiac arrest without STE/cardio. shock O GRACE >140
	H3		
	%rising		

Progress	Order for one day	Order for continuation
____/____/____	O Admit O CCU O CBC, PT, PTT, INR O BUN, Cr, E'lyte, Ca, Mg, P O Anti-HIV O hs-troponin I Hr.0 and 3 ถ้าเจ้าจาก ราช. และไม่ต้องเจาะช้ำ	O Regular diet O Record V/S, I/O
____:____		Medication
Pre-coun. anti_HIV	O CXR O EKG 12 leads O NSS 500 ml IV 40 ml/hr Lt. arm if AVF, on IV at non-AVF arm O on O2 canula 3 L if O2sat <90% O ASA (300) 1 tab chew stat O Clopidogrel GPO (75) 4 tabs PO stat O ISDN (5) 1 tab SL prn chest pain	O ASA (81) 1x1 pc O Clopidogrel GPO (75) 1x1 pc O Omeprazole (20) 1x1 ac O Atorvastatin (40) 1x1 hs O Senokot 2x1 hs O Ativan (0.5) 1x1 hs O ISDN (5) 1 tab SL prn chest pain
	(ER/ward sign)_____	O Fondaparinux
	O Standby Echo IPD case O Standby CAG O พรุน FBS, lipid O Consult cardiac rehab O Consult เลิกบุหรี่ (ถ้ายังสูบบุหรี่)	O 2.5 mg SC OD x 5 Days (ห้ามให้ถ้า GFR <20)
		O Enoxaparin
	(ward sign)_____	O _____ ml SC q _____ hr x 5 Days Dose of enoxaparin - Age<75; 1 mg/kg SC q 12 hr - Age≥75; 0.75 mg/kg SC q 12 hr - If GFR 15-30; give OD dose
		O Heparin
Consult		ใช้ในกรณี GFR <15 O _____ u (60 u/kg) IV bolus then heparin 10000 u + NSS 100 ml IV drip _____ u/hr (12 u/kg/hr) MAX 1000 u/hr x 2 Days
- Rehab ____/____/____	(ward sign)_____	(ward sign)_____
- Smoke ____/____/____		

Ver 2 031164